



SRPK2 rabbit pAb

Cat No.:ES10216

For research use only

Overview

Product Name	SRPK2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SRPK2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C . Avoid repeated freeze-thaw cycles.
Protein Name	SRSF protein kinase 2 (EC 2.7.11.1) (SFRS protein kinase 2) (Serine/arginine-rich protein-specific kinase 2) (SR-protein-specific kinase 2) [Cleaved into: SRSF protein kinase 2 N-terminal; SRSF protei
Gene Name	SRPK2
Cellular localization	Cytoplasm . Nucleus, nucleoplasm . Nucleus speckle . Chromosome . Shuttles between the nucleus and the cytoplasm (PubMed:19592491, PubMed:21157427, PubMed:21056976). KAT5/TIP60 inhibits its nuclear translocation (PubMed:21157427). Phosphorylation at Thr-492 by PKB/AKT1 promotes nuclear translocation (PubMed:19592491). Preferentially localizes across the entire gene coding region (PubMed:28076779). During transcription, accumulates at chromatin loci where unscheduled R-loops form and colocalizes with paused 'Ser-5'-phosphorlyated POLR2A/RNA polymerase II and helicase DDX23 (PubMed:28076779). .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using





ELK Biotechnology

Clonality

epitope-specific immunogen.

Concentration

Polyclonal

Observed band

1 mg/ml

Human Gene ID

75kD

Human Swiss-Prot Number

6733

Alternative Names

P78362

Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Ser-52 and Ser-588.,function:Phosphorylates RS domain-containing proteins, such as SFRS1 and SFRS2 on serine residues. Role in spliceosome assembly and in mediating the trafficking of splicing factors. Appears to mediate HBV core protein phosphorylation which is a prerequisite for pregenomic RNA encapsidation into viral capsids.,sequence caution:The cDNA appears to contain a duplicated region.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,tissue specificity:Highly expressed in brain, moderately expressed in heart and skeletal muscle and at low levels in lung, liver, and kidney.,



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C