

BAP1 rabbit pAb

Cat No.: ES10425

For research use only

Overview

Product Name BAP1 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein . at

AA range: 150-230

Specificity BAP1 Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

3.4.19.12) (BRCA1-associated protein 1) (Cerebral

protein 6)

Gene Name BAP1 KIAA0272 hucep-6

Cellular localization Cytoplasm . Nucleus . Mainly nuclear. Binds to

chromatin. Localizes to the cytoplasm when monoubiquitinated by the E2/E3 hybrid

ubiquitin-protein ligase UBE2O

(PubMed:24703950). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 80kD
Human Gene ID 8314
Human Swiss-Prot Number 092560

Alternative Names

Background This gene belongs to the ubiquitin C-terminal

hydrolase subfamily of deubiquitinating enzymes that are involved in the removal of ubiquitin from



+86-27-59760950 ELKbio@ELKbiotech.com ww



proteins. The encoded enzyme binds to the breast cancer type 1 susceptibility protein (BRCA1) via the RING finger domain of the latter and acts as a tumor suppressor. In addition, the enzyme may be involved in regulation of transcription, regulation of cell cycle and growth, response to DNA damage and chromatin dynamics. Germline mutations in this gene may be associated with tumor predisposition syndrome (TPDS), which involves increased risk of cancers including malignant mesothelioma, uveal melanoma and cutaneous melanoma. [provided by RefSeq, May 2013],



+86-27-59760950