

CAD12 rabbit pAb

Cat No.:ES10503

For research use only

Overview

Product Name	CAD12 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at
	AA range: 500-580
Specificity	CAD12 Polyclonal Antibody detects endogenous
	levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles.
Protein Name	Cadherin-12 (Brain cadherin) (BR-cadherin) (Neural
	type cadherin 2) (N-cadherin 2)
Gene Name	CDH12
Cellular localization	Cell membrane; Single-pass type I membrane
	protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	87kD
Human Gene ID	1010
Human Swiss-Prot Number	P55289
Alternative Names	
Background	This gene encodes a type II classical cadherin of the
	cadherin superfamily. Alternative splicing of this
	gene results in multiple transcript variants. At least
	one of these variants encodes a preproprotein that
	is proteolytically processed to generate the mature
	cadherin protein. These integral membrane proteins
	mediate calcium-dependent cell-cell adhesion and



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



are composed of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. Type II (atypical) cadherins are defined based on their lack of a histidine-alanine-valine (HAV) cell adhesion recognition sequence specific to type I cadherins. This particular cadherin appears to be expressed specifically in the brain and its temporal pattern of expression would be consistent with a role during a critical period of neuronal development, perhaps specifically during synaptogenesis. [p



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C