

CIC rabbit pAb

Cat No.: ES10630

For research use only

Overview

Product Name CIC rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein . at

AA range: 550-630

Specificity CIC Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20° C. Avoid repeated freeze-thaw cycles.

Protein Name Protein capicua homolog

Gene Name CIC KIAA0306
Cellular localization Nucleus .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 176kD
Human Gene ID 23152
Human Swiss-Prot Number Q96RK0

Alternative Names

Background The protein encoded by this gene is an ortholog of

the Drosophila melanogaster capicua gene, and is a member of the high mobility group (HMG)-box superfamily of transcriptional repressors. This protein contains a conserved HMG domain that is involved in DNA binding and nuclear localization, and a conserved C-terminus. Studies suggest that the N-terminal region of this protein interacts with

Atxn1 (GeneID:6310), to form a transcription



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com



repressor complex, and in vitro studies suggest that polyglutamine-expansion of ATXN1 may alter the repressor activity of this complex. Mutations in this gene have been associated with olidogdendrogliomas (PMID:21817013). In addition, translocation events resulting in gene fusions of this gene with both DUX4 (GeneID:100288687) and FOXO4 (GeneID:4303) have been associated with round cell sarcomas. There are multiple pseudogenes of this gene found on chromosomes



+86-27-59760950