



# S35C2 rabbit pAb

Cat No.:ES10656

For research use only

## Overview

<b>Product Name</b>	S35C2 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Rat;Mouse;
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	S35C2 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Solute carrier family 35 member C2 (Ovarian cancer-overexpressed gene 1 protein)
<b>Gene Name</b>	SLC35C2 C20orf5 OVCOV1 CGI-15
<b>Cellular localization</b>	Membrane ; Multi-pass membrane protein . Golgi apparatus, cis-Golgi network membrane . Endoplasmic reticulum-Golgi intermediate compartment membrane .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	40kD
<b>Human Gene ID</b>	51006
<b>Human Swiss-Prot Number</b>	Q9NQQ7
<b>Alternative Names</b>	
<b>Background</b>	This gene encodes a member of the triose-phosphate transporter protein family. This gene is regulated by oxygen tension, is induced in hypoxic trophoblast cells, and is overexpressed in ovarian cancer. Alternative splicing results in





multiple transcript variants. A pseudogene of this gene has been defined on the X chromosome. [provided by RefSeq, Jul 2013],

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night

