

## GAR1 rabbit pAb

Cat No.: ES10659

For research use only

## Overview

Product Name GAR1 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

**Immunogen** Synthesized peptide derived from human protein .

at AA range: 110-190

**Specificity** GAR1 Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name H/ACA ribonucleoprotein complex subunit 1

(Nucleolar protein family A member 1) (snoRNP

protein GAR1)

Gene Name GAR1 NOLA1

**Cellular localization** Nucleus, nucleolus. Nucleus, Cajal body. Also

localized to Cajal bodies (coiled bodies).

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 23kD
Human Gene ID 54433
Human Swiss-Prot Number Q9NY12

**Alternative Names** 

**Background** This gene is a member of the H/ACA snoRNPs (small

nucleolar ribonucleoproteins) gene family. snoRNPs are involved in various aspects of rRNA processing and modification and have been classified into two families: C/D and H/ACA. The H/ACA snoRNPs also include the DKC1, NOLA2 and NOLA3 proteins.




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



These four H/ACA snoRNP proteins localize to the dense fibrillar components of nucleoli and to coiled (Cajal) bodies in the nucleus. Both 18S rRNA production and rRNA pseudouridylation are impaired if any one of the four proteins is depleted. These four H/ACA snoRNP proteins are also components of the telomerase complex. The encoded protein of this gene contains two glycine-and arginine-rich domains and is related to Saccharomyces cerevisiae Gar1p. Two splice variants have been found for this gene. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

