

## CR3L3 rabbit pAb

Cat No.: ES10755

For research use only

## Overview

Product Name CR3L3 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from part region of

human protein AA range: 1-100

**Specificity** CR3L3 Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Cyclic AMP-responsive element-binding protein

3-like protein 3 (cAMP-responsive element-binding

protein 3-like protein 3) (Transcription factor CREB-H) [Cleaved into: Processed cyclic

AMP-responsive e

Gene Name CREB3L3 CREBH HYST1481

**Cellular localization** Endoplasmic reticulum membrane; Single-pass type

II membrane protein .; [Processed cyclic

AMP-responsive element-binding protein 3-like protein 3]: Nucleus . Under ER stress the cleaved N-terminal cytoplasmic domain translocates into the

nucleus.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 50kD
Human Gene ID 84699
Human Swiss-Prot Number Q68CJ9

**Alternative Names** 



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



**Background** 

This gene encodes a member of the basic-leucine zipper family and the AMP-dependent transcription factor family. The encoded protein is localized to the endoplasmic reticulum and acts as a transcription factor activated by cyclic AMP stimulation. The encoded protein binds the cyclic AMP response element (CRE) and the box-B element and has been linked to acute inflammatory response, hepatocellular carcinoma, triglyceride metabolism, and hepcidin expression. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2012],



+86-27-59760950