

KPCD3 rabbit pAb

Cat No.: ES10822

For research use only

Overview

Product Name KPCD3 rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from part region of

human protein

Specificity KPCD3 Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

StorageStore at -20 °C. Avoid repeated freeze-thaw cycles.Protein NameSerine/threonine-protein kinase D3 (EC 2.7.11.13)

(Protein kinase C nu type) (Protein kinase EPK2)

(nPKC-nu)

Gene Name PRKD3 EPK2 PRKCN

Cellular localization Cytoplasm . Membrane . Translocation to the cell

membrane is required for kinase activation.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band97kDHuman Gene ID23683Human Swiss-Prot Number094806

Alternative Names

Background This gene belongs to the multigene protein kinase D

family of serine/threonine kinases, which bind diacylglycerol and phorbol esters. Members of this family are characterized by an N-terminal regulatory

domain comprised of a tandem repeat of cysteine-rich zinc-finger motifs and a pleckstrin



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



domain. The C-terminal region contains the catalytic domain and is distantly related to calcium-regulated kinases. Catalytic activity of this enzyme promotes its nuclear localization. This protein has been implicated in a variety of functions including negative regulation of human airway epithelial barrier formation, growth regulation of breast and prostate cancer cells, and vesicle trafficking. [provided by RefSeq, Jan 2015],



+86-27-59760950

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C