

MXI1 rabbit pAb

Cat No.: ES11015

For research use only

Overview

Product Name MXI1 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from part region of

human protein

Specificity MXI1 Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Max-interacting protein 1 (Max interactor 1) (Class C

basic helix-loop-helix protein 11) (bHLHc11)

Gene Name MXI1 BHLHC11

Cellular localization Nucleus.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 25kD
Human Gene ID 4601
Human Swiss-Prot Number P50539

Alternative Names

Background Expression of the c-myc gene, which produces an

oncogenic transcription factor, is tightly regulated in normal cells but is frequently deregulated in human cancers. The protein encoded by this gene is a transcriptional repressor thought to negatively regulate MYC function, and is therefore a potential


tumor suppressor. This protein inhibits the transcriptional activity of MYC by competing for



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





MAX, another basic helix-loop-helix protein that binds to MYC and is required for its function. Defects in this gene are frequently found in patients with prostate tumors. Three alternatively spliced transcripts encoding different isoforms have been described. Additional alternatively spliced transcripts may exist but the products of these transcripts have not been verified experimentally. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night

