

## **PTPRM** rabbit pAb

## Cat No.:ES11058

For research use only

## Overview

Product Name	PTPRM rabbit pAb	
Host species	Rabbit	
Applications	WB;ELISA	
Species Cross-Reactivity	Human;Mouse	
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000	
Immunogen	Synthesized peptide derived from part region of	
	human protein. AA range 21-61	
Specificity	PTPRM Polyclonal Antibody detects endogenous	
	levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze-thaw cycles.	
Protein Name	Receptor-type tyrosine-protein phosphatase mu	
	(Protein-tyrosine phosphatase mu) (R-PTP-mu) (EC	
	3.1.3.48)	
Gene Name	PTPRM PTPRL1	
Cellular localization	Cell membrane ; Single-pass type I membrane	
	protein. Localizes in regions of cell-cell contact	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	159kD	
Human Gene ID	5797	
Human Swiss-Prot Number	P28827	
Alternative Names		
Background	The protein encoded by this gene is a member of	
	the protein tyrosine phosphatase (PTP) family. PTPs	
	are known to be signaling molecules that regulate a	
	variety of cellular processes including cell growth,	
	differentiation, mitotic cycle, and oncogenic	
	transformation. This PTP possesses an extracellular	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



region, a single transmembrane region, and two tandem catalytic domains, and thus represents a receptor-type PTP. The extracellular region contains a meprin-A5 antigen-PTP mu (MAM) domain, an Ig-like domain and four fibronectin type III-like repeats. This PTP has been shown to mediate cell-cell aggregation through the interaction with another molecule of this PTP on an adjacent cell. This PTP can interact with scaffolding protein RACK1/GNB2L1, which may be necessary for the downstream signaling in response to cell-cell adhesion. Alternative splicing results in multiple transcrip



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C