

TCAM2 rabbit pAb

Cat No.:ES11113

For research use only

Overview

Product Name	TCAM2 rabbit pAb	
Host species	Rabbit	
Applications	WB;ELISA	
Species Cross-Reactivity	Human;Mouse	
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000	
Immunogen	Synthesized peptide derived from part region of	
-	human protein	
Specificity	TCAM2 Polyclonal Antibody detects endogenous	
	levels of protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	TIR domain-containing adapter molecule 2	
	(TICAM-2) (Putative NF-kappa-B-activating protein	
	502) (TRIF-related adapter molecule) (Toll-like	
	receptor adaptor protein 3) (Toll/interleukin-1	
	receptor doma	
Gene Name	TICAM2 TIRAP3 TIRP TRAM	
Cellular localization	[Isoform 1]: Cytoplasm . Golgi apparatus. Cell	
	membrane . Endoplasmic reticulum. Early	
	endosome membrane. Late endosome membrane.	
	Cell projection, phagocytic cup . Localized to the	
	plasma membrane as a result of myristoylation.	
	Phosphorylation on Ser-16 l	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	25kD	
Human Gene ID	100302736	
Human Swiss-Prot Number	Q86XR7	
Alternative Names		



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

TIRP is a Toll/interleukin-1 receptor (IL1R; MIM 147810) (TIR) domain-containing adaptor protein involved in Toll receptor signaling (see TLR4; MIM 603030).[supplied by OMIM, Apr 2004],

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C