

BGH3 rabbit pAb

Cat No.: ES11167

For research use only

Overview

Product Name BGH3 rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein . at

AA range: 230-310

Specificity BGH3 Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20 °C. Avoid repeated freeze-thaw cycles.

Protein Name Transforming growth factor-beta-induced protein

is 1-2 (Posterio 1-2) ((Complete a sixth all in) (PCP) and sixth all in) (PCP).

ig-h3 (Beta ig-h3) (Kerato-epithelin) (RGD-containing

collagen-associated protein) (RGD-CAP)

Gene Name TGFBI BIGH3

Cellular localization Secreted . Secreted, extracellular space, extracellular

matrix . May be associated both with microfibrils and with the cell surface (PubMed:8077289). .
The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 75kD
Human Gene ID 7045
Human Swiss-Prot Number Q15582

Alternative Names

Purification

Background This gene encodes an RGD-containing protein that

binds to type I, II and IV collagens. The RGD motif is

found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



protein plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy. [provided by RefSeq, Jul 2008],



+86-27-59760950