

OLR1 rabbit pAb

Cat No.: ES11225

For research use only

Overview

Product Name OLR1 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein .

at AA range: 60-140

Specificity OLR1 Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Oxidized low-density lipoprotein receptor 1 (Ox-LD

in Name
Oxidized low-density lipoprotein receptor 1 (Ox-LDL receptor 1) (C-type lectin domain family 8 member
A) (Lectin-like oxidized LDL receptor 1) (LOX-1)

(Lectin-like oxLDL receptor 1) (hLOX-1) (Lectin-t

Gene Name OLR1 CLEC8A LOX1

Cellular localization Cell membrane; Lipid-anchor. Cell membrane;

Single-pass type II membrane protein. Membrane

raft. Secreted. A secreted form also exists. Localization to membrane rafts requires

palmitoylation.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 30kD
Human Gene ID 4973
Human Swiss-Prot Number P78380

Alternative Names

Background This gene encodes a low density lipoprotein

receptor that belongs to the C-type lectin




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



superfamily. This gene is regulated through the cyclic AMP signaling pathway. The encoded protein binds, internalizes and degrades oxidized low-density lipoprotein. This protein may be involved in the regulation of Fas-induced apoptosis. This protein may play a role as a scavenger receptor. Mutations of this gene have been associated with atherosclerosis, risk of myocardial infarction, and may modify the risk of Alzheimer's disease. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Feb 2010],

Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950