

## **RYK rabbit pAb**

## Cat No.:ES11257

For research use only

## Overview

Product Name	RYK rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at
U U	AA range: 150-230
Specificity	RYK Polyclonal Antibody detects endogenous levels
	of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!{ m C}$ . Avoid repeated freeze-thaw cycles.
Protein Name	Tyrosine-protein kinase RYK (EC 2.7.10.1)
Gene Name	RYK JTK5A
Cellular localization	Membrane ; Single-pass type I membrane protein .
	Nucleus . Cytoplasm . In cells that have undergone
	neuronal differentiation, the C-terminal cleaved part
	is translocated from the cytoplasm to the nucleus
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	66kD
Human Gene ID	6259
Human Swiss-Prot Number	P34925
Alternative Names	
Background	The protein encoded by this gene is an atypical
	member of the family of growth factor receptor
	protein tyrosine kinases, differing from other
	members at a number of conserved residues in the
	activation and nucleotide binding domains. This
	gene product belongs to a subfamily whose



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



members do not appear to be regulated by phosphorylation in the activation segment. It has been suggested that mediation of biological activity by recruitment of a signaling-competent auxiliary protein may occur through an as yet uncharacterized mechanism. The encoded protein has a leucine-rich extracellular domain with a WIF-type Wnt binding region, a single transmembrane domain, and an intracellular tyrosine kinase domain. This protein is involved in stimulating Wnt signaling pathways such as the regulation of axon pathfinding. Alternative splicing results in multiple transcript variants encoding distinct isoforms



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C