

PLOD2 rabbit pAb

Cat No.:ES11324

For research use only

Overview

| Product Name | PLOD2 rabbit pAb |
|--------------------------|--|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human; Rat; Mouse |
| Recommended dilutions | WB 1:500-2000 ELISA 1:5000-20000 |
| Immunogen | Synthesized peptide derived from human protein . at |
| <u> </u> | AA range: 600-680 |
| Specificity | PLOD2 Polyclonal Antibody detects endogenous |
| | levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles. |
| Protein Name | Procollagen-lysine,2-oxoglutarate 5-dioxygenase 2 |
| | (EC 1.14.11.4) (Lysyl hydroxylase 2) (LH2) |
| Gene Name | PLOD2 |
| Cellular localization | Rough endoplasmic reticulum membrane; Peripheral |
| | membrane protein; Lumenal side. |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 81kD |
| Human Gene ID | 5352 |
| Human Swiss-Prot Number | O00469 |
| Alternative Names | |
| Background | The protein encoded by this gene is a |
| | membrane-bound homodimeric enzyme that is |
| | localized to the cisternae of the rough endoplasmic |
| | reticulum. The enzyme (cofactors iron and |
| | ascorbate) catalyzes the hydroxylation of lysyl |
| | residues in collagen-like peptides. The resultant |
| | hydroxylysyl groups are attachment sites for |
| | |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



carbohydrates in collagen and thus are critical for the stability of intermolecular crosslinks. Some patients with Ehlers-Danlos syndrome type VIB have deficiencies in lysyl hydroxylase activity. Mutations in the coding region of this gene are associated with Bruck syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C