

O51M1 rabbit pAb

Cat No.:ES11520

For research use only

Overview

| Product Name | O51M1 rabbit pAb |
|------------------------------|---|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | WB 1:500-2000 ELISA 1:5000-20000 |
| Immunogen | Synthesized peptide derived from part region of |
| | human protein |
| Specificity | O51M1 Polyclonal Antibody detects endogenous |
| | levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Olfactory receptor 51M1 (Odorant receptor |
| | HOR5'beta7) (Olfactory receptor OR11-40) |
| Gene Name | OR51M1 |
| Cellular localization | Cell membrane; Multi-pass membrane protein. |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 34kD |
| Human Gene ID | |
| Human Swiss-Prot Number | Q9H341 |
| Alternative Names | |
| Background | Olfactory receptors interact with odorant molecules |
| | in the nose, to initiate a neuronal response that |
| | triggers the perception of a smell. The olfactory |
| | receptor proteins are members of a large family of |
| | G-protein-coupled receptors (GPCR) arising from |
| | single coding-exon genes. Olfactory receptors share |
| | a 7-transmembrane domain structure with many |
| | |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


neurotransmitter and hormone receptors and are

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C