

OR8D4 rabbit pAb

Cat No.: ES11614

For research use only

Overview

Product Name OR8D4 rabbit pAb

Host species Rabbit WB;ELISA **Applications**

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein.

at AA range: 30-110

OR8D4 Polyclonal Antibody detects endogenous Specificity

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Olfactory receptor 8D4 (Olfactory receptor **Protein Name**

OR11-275)

OR8D4 **Gene Name**

Cellular localization Cell membrane; Multi-pass membrane protein. Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml Observed band 34kD **Human Gene ID** 338662 **Human Swiss-Prot Number** Q8NGM9

Alternative Names

Background olfactory receptor family 8 subfamily D member

4(OR8D4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of


G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com