

KLC1 rabbit pAb

Cat No.: ES11766

For research use only

Overview

Product Name KLC1 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from part region of

human protein AA range: 128-178

Specificity KLC1 Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20° C. Avoid repeated freeze-thaw cycles.

Protein Name Kinesin light chain 1 (KLC 1)

Gene Name KLC1 KLC KNS2

Cellular localization Cell projection, growth cone . Cytoplasmic vesicle .

Cytoplasm, cytoskeleton.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 63kD
Human Gene ID 3831
Human Swiss-Prot Number Q07866

Alternative Names

Background Conventional kinesin is a tetrameric molecule

composed of two heavy chains and two light chains, and transports various cargos along microtubules toward their plus ends. The heavy chains provide the motor activity, while the light chains bind to various cargos. This gene encodes a member of the kinesin light chain family. It associates with kinesin heavy chain through an N-terminal domain, and six



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



tetratricopeptide repeat (TPR) motifs are thought to be involved in binding of cargos such as vesicles, mitochondria, and the Golgi complex. Thus, kinesin light chains function as adapter molecules and not motors per se. Although previously named "kinesin 2", this gene is not a member of the kinesin-2 / kinesin heavy chain subfamily of kinesin motor proteins. Extensive alternative splicing produces isoforms with different C-termini that are proposed to bind to different cargos; however, the full-



+86-27-59760950