



# HIC2 rabbit pAb

Cat No.:ES11909

For research use only

## Overview

<b>Product Name</b>	HIC2 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	HIC2 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Hypermethylated in cancer 2 protein (Hic-2) (Hic1-related gene on chromosome 22 protein) (Hic-3) (Zinc finger and BTB domain-containing protein 30)
<b>Gene Name</b>	HIC2 HRG22 KIAA1020 ZBTB30
<b>Cellular localization</b>	Nucleus.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	67kD
<b>Human Gene ID</b>	23119
<b>Human Swiss-Prot Number</b>	Q96JB3
<b>Alternative Names</b>	
<b>Background</b>	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,

