



HDAC5 (phospho Ser498) rabbit pAb

Cat No.:ES1324

For research use only

Overview

Product Name	HDAC5 (phospho Ser498) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human HDAC5 around the phosphorylation site of Ser498. AA range:464-513
Specificity	Phospho-HDAC5 (S498) Polyclonal Antibody detects endogenous levels of HDAC5 protein only when phosphorylated at S498.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Histone deacetylase 5
Gene Name	HDAC5
Cellular localization	Nucleus. Cytoplasm. Shuttles between the nucleus and the cytoplasm. In muscle cells, it shuttles into the cytoplasm during myocyte differentiation. The export to cytoplasm depends on the interaction with a 14-3-3 chaperone protein and is due to its phosphorylation at Ser-259 and Ser-498 by AMPK, CaMK1 and SIK1.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	122kD
Human Gene ID	10014





Human Swiss-Prot Number


Q9UQL6

Alternative Names


HDAC5; KIAA0600; Histone deacetylase 5; HD5; Antigen NY-CO-9

Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using Phospho-HDAC5 (S498) Polyclonal Antibody diluted at 1:500





Western Blot analysis of VEC cells using Phospho-HDAC5 (S498) Polyclonal Antibody diluted at 1:500





Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using HDAC5 (Phospho-Ser498) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells, using HDAC5 (Phospho-Ser498) Antibody. The lane on the right is blocked with the phospho peptide.

