

IκB-β (phospho Ser23) rabbit pAb

Cat No.:ES1346

For research use only

Overview

Product Name	ΙκΒ-β (phospho Ser23) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
-	peptide derived from human IkappaB-beta around
	the phosphorylation site of Ser23. AA range:8-57
Specificity	Phospho-IκB- β (S23) Polyclonal Antibody detects
	endogenous levels of ΙκΒ-β protein only when
	phosphorylated at S23.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	NF-kappa-B inhibitor beta
Gene Name	NFKBIB
Cellular localization	Cytoplasm . Nucleus .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	37kD
Human Gene ID	4793
Human Swiss-Prot Number	Q15653
Alternative Names	NFKBIB; IKBB; TRIP9; NF-kappa-B inhibitor beta;
	NF-kappa-BIB; I-kappa-B-beta; IkB-B; IkB-beta;
	IkappaBbeta; Thyroid receptor-interacting protein 9;
	TR-interacting protein 9; TRIP-9
Background	The protein encoded by this gene belongs to the



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



NF-kappa-B inhibitor family, which inhibit NF-kappa-B by complexing with, and trapping it in the cytoplasm. Phosphorylation of serine residues on these proteins by kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B, which translocates to the nucleus to function as a transcription factor. Alternatively spliced transcript variants have been found for this gene.[provided by RefSeq, Jul 2011],

Western Blot analysis of various cells using Phospho-IκB-β (S23) Polyclonal Antibody



HeLa

(kD)

117-85-

48-

34-

26-

19-

Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IkappaB-beta (Phospho-Ser23) Antibody




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Immunofluorescence analysis of HeLa cells treated with TNF-a 20nM 15', using IkappaB-beta (Phospho-Ser23) Antibody. The picture on the right is blocked with the phospho peptide.

Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using IkappaB-beta (Phospho-Ser23) Antibody. The picture on the right is blocked with the phospho peptide.





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C