

Paxillin (phospho Tyr118) rabbit pAb

Cat No.: ES1392

For research use only

Overview

Specificity

Product Name Paxillin (phospho Tyr118) rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human Paxillin around the phosphorylation site of Tyr118. AA range:85-134 Phospho-Paxillin (Y118) Polyclonal Antibody detects

endogenous levels of Paxillin protein only when

phosphorylated at Y118.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Paxillin Gene Name PXN

Cellular localization Cytoplasm, cytoskeleton . Cell junction, focal

adhesion . Cytoplasm, cell cortex . Colocalizes with integrins at the cell periphery. Colocalize with PXN to membrane ruffles and the leading edge of

migrating cells (PubMed:23128389). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 64kD
Human Gene ID 5829
Human Swiss-Prot Number P49023
Alternative Names PXN; Paxillin

Background This gene encodes a cytoskeletal protein involved in

actin-membrane attachment at sites of cell adhesion




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



to the extracellular matrix (focal adhesion). Alternatively spliced transcript variants encoding different isoforms have been described for this gene. These isoforms exhibit different expression pattern, and have different biochemical, as well as physiological properties (PMID:9054445). [provided by RefSeq, Aug 2011],

Western Blot analysis of various cells using Phospho-Paxillin (Y118) Polyclonal Antibody



-- 117
-- 85

PAXILLIN/PAXI -- (pTyr118)
-- 48
-- 34
-- 26
-- 19 (kD)

Western blot analysis of lysates from HeLa cells treated with EGF, using Paxillin (Phospho-Tyr118) Antibody. The lane on the right is blocked with the phospho peptide.

