

Amylin rabbit pAb

Cat No.: ES1651

For research use only

Overview

Product Name Amylin rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human Amylin. AA range:1-50

Specificity Amylin Polyclonal Antibody detects endogenous

levels of Amylin protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Islet amyloid polypeptide

Gene Name IAPP
Cellular localization Secreted .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 25kD
Human Gene ID 3375
Human Swiss-Prot Number P10997

Alternative Names IAPP; Islet amyloid polypeptide; Amylin;

Diabetes-associated peptide; DAP; Insulinoma

amyloid peptide

Background islet amyloid polypeptide(IAPP) Homo sapiens

This gene encodes a member of the calcitonin family of peptide hormones. This hormone is released from

pancreatic beta cells following food intake to



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




regulate blood glucose levels and act as a satiation signal. Human patients with type 1 and advanced type 2 diabetes exhibit reduced levels of the encoded hormone in blood and pancreas. This protein also exhibits a bactericidal, antimicrobial activity. [provided by RefSeq, Jul 2016],

Western Blot analysis of various cells using Amylin Polyclonal Antibody diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Amylin Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, treated with EGF 200ng/ml 30', using Amylin Antibody. The lane on the right is blocked with the synthesized peptide.

