

## **BMP-7** rabbit pAb

## Cat No.:ES1779

For research use only

## Overview

Product Name	BMP-7 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human; Mouse; Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human BMP-7. AA
	range:124-173
Specificity	BMP-7 Polyclonal Antibody detects endogenous
	levels of BMP-7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Bone morphogenetic protein 7
Gene Name	BMP7
Cellular localization	Secreted.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	55kD
Human Gene ID	655
Human Swiss-Prot Number	P18075
Alternative Names	BMP7; OP1; Bone morphogenetic protein 7; BMP-7;
	Osteogenic protein 1; OP-1; Eptotermin alfa
Background	This gene encodes a secreted ligand of the TGF-beta
	(transforming growth factor-beta) superfamily of
	proteins. Ligands of this family bind various
	TGF-beta receptors leading to recruitment and
	activation of SMAD family transcription factors that



+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer, which plays a role in bone, kidney and brown adipose tissue development. Additionally, this protein induces ectopic bone formation and may promote fracture healing in human patients. [provided by RefSeq, Jul 2016],



133----100---55---40---25----15---- Western Blot analysis of various cells using BMP-7 Polyclonal Antibody diluted at 1:500

Western Blot analysis of rat-musle cells using BMP-7 Polyclonal Antibody diluted at 1:500

Immunohistochemistry analysis of BMP-7 antibody in paraffin-embedded human tonsil tissue.






+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C.





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C