

## Cyclin E1 rabbit pAb

## Cat No.:ES2082

For research use only

## Overview

Product Name	Cyclin E1 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat;Pig
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human Cyclin E1. AA
	range:91-140
Specificity	Cyclin E1 Polyclonal Antibody detects endogenous
	levels of Cyclin E1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze-thaw cycles.
Protein Name	G1/S-specific cyclin-E1
Gene Name	CCNE1
Cellular localization	Nucleus .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	49kD
Human Gene ID	898
Human Swiss-Prot Number	P24864
Alternative Names	CCNE1; CCNE; G1/S-specific cyclin-E1
Background	The protein encoded by this gene belongs to the
	highly conserved cyclin family, whose members are
	characterized by a dramatic periodicity in protein
	abundance through the cell cycle. Cyclins function as
	regulators of CDK kinases. Different cyclins exhibit



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C