

Di-Ras1 rabbit pAb

Cat No.:ES2174

For research use only

Overview

Product Name	Di-Ras1 rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human;Rat;Mouse;	
Recommended dilutions	Western Blot: 1/500 - 1/2000.	
	Immunohistochemistry: 1/100 - 1/300.	
	Immunofluorescence: 1/200 - 1/1000. ELISA:	
	1/20000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
-	peptide derived from human DIRA1. AA	
	range:149-198	
Specificity	Di-Ras1 Polyclonal Antibody detects endogenous	
	levels of Di-Ras1 protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	GTP-binding protein Di-Ras1	
Gene Name	DIRAS1	
Cellular localization	Cell membrane ; Lipid-anchor ; Cytoplasmic side .	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	22kD	
Human Gene ID	148252	
Human Swiss-Prot Number	O95057	
Alternative Names	DIRAS1; GBTS1; RIG; GTP-binding protein Di-Ras1;	
	Distinct subgroup of the Ras family member 1;	
	Ras-related inhibitor of cell growth; Rig; Small	
	GTP-binding tumor suppressor 1	
Background	DIRAS1 belongs to a distinct branch of the	
	functionally diverse Ras (see HRAS; MIM 190020)	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



HeLa

(kD)

117-85-

48-

34-

26-

19-

DIRA1 -

HeLa HepG2

-- 117

-- 85


-- 48 -- 34

-- 26

-- 19 (kD) superfamily of monomeric GTPases.[supplied by OMIM, Apr 2004],

Western Blot analysis of various cells using Di-Ras1 Polyclonal Antibody

Western blot analysis of lysates from HeLa and HepG2 cells, using DIRA1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C