

## Follistatin rabbit pAb

**Cat No.:ES2357** 

For research use only

## Overview

Product Name Follistatin rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human FST. AA range:121-170

**Specificity** Follistatin Polyclonal Antibody detects endogenous

levels of Follistatin protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Follistatin

Gene Name FST

Cellular localization Secreted.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 65kD
Human Gene ID 10468
Human Swiss-Prot Number P19883

Human Swiss-Prot Number P19883
Alternative Names FST; Follows

Alternative Names FST; Follistatin; FS; Activin-binding protein Background Follistatin is a single-chain gonadal protein that

specifically inhibits follicle-stimulating hormone release. The single FST gene encodes two isoforms, FST317 and FST344 containing 317 and 344 amino acids respectively, resulting from alternative splicing of the precursor mRNA. In a study in which 37 candidate genes were tested for

+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com





linkage and association with polycystic ovary syndrome (PCOS) or hyperandrogenemia in 150 families, evidence was found for linkage between PCOS and follistatin. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Follistatin Polyclonal Antibody diluted at 1:500



HepG2


(kD)

117-85-

48-

34-

Western blot analysis of the lysates from COLO205 cells using FST antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

