

GRB10 rabbit pAb

Cat No.:ES2472

For research use only

Overview

Product Name GRB10 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

The antiserum was produced against synthesize.

Immunogen The antiserum was produced against synthesized

peptide derived from human GRB10. AA

range:33-82

Specificity GRB10 Polyclonal Antibody detects endogenous

levels of GRB10 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Growth factor receptor-bound protein 10

Gene Name GRB10

Cellular localization Cytoplasm . When complexed with NEDD4 and

IGF1R, follows IGF1R internalization, remaining associated with early endosomes. Uncouples from IGF1R-containing endosomes before the sorting of the receptor to the lysosomal compartment (By

similarity). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 67kD
Human Gene ID 2887
Human Swiss-Prot Number O13322

Alternative Names GRB10; GRBIR; KIAA0207; Growth factor






Background

receptor-bound protein 10; GRB10 adapter protein; Insulin receptor-binding protein Grb-IR The product of this gene belongs to a small family of adapter proteins that are known to interact with a number of receptor tyrosine kinases and signaling molecules. This gene encodes a growth factor receptor-binding protein that interacts with insulin receptors and insulin-like growth-factor receptors. Overexpression of some isoforms of the encoded protein inhibits tyrosine kinase activity and results in growth suppression. This gene is imprinted in a highly isoform- and tissue-specific manner, with expression observed from the paternal allele in the brain, and from the maternal allele in the placental trophoblasts. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2010],

Western Blot analysis of various cells using GRB10 Polyclonal Antibody diluted at 1:2000



27

Western Blot analysis of 3T3 cells using GRB10 Polyclonal Antibody diluted at 1:2000





+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com







Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GRB10 Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950

