

## Histamine H1 Receptor rabbit pAb

Cat No.: ES2516

For research use only

## Overview

Product Name Histamine H1 Receptor rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human HRH1. AA

range:141-190

**Specificity** Histamine H1 Receptor Polyclonal Antibody detects

endogenous levels of Histamine H1 Receptor

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Histamine H1 receptor

Gene Name HRH1

Cellular localizationCell membrane ; Multi-pass membrane protein .PurificationThe antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 60kD
Human Gene ID 3269
Human Swiss-Prot Number P35367

Alternative Names HRH1; Histamine H1 receptor; H1R; HH1R

Background Histamine is a ubiquitous messenger molecule

released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. The protein encoded by this gene is an integral




+86-27-59760950 ELKbio@ELKbiotech.com w



membrane protein and belongs to the G protein-coupled receptor superfamily. It mediates the contraction of smooth muscles, the increase in capillary permeability due to contraction of terminal venules, the release of catecholamine from adrenal medulla, and neurotransmission in the central nervous system. It has been associated with multiple processes, including memory and learning, circadian rhythm, and thermoregulation. It is also known to contribute to the pathophysiology of allergic diseases such as atopic dermatitis, asthma, anaphylaxis and allergic rhinitis. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by Ref

(kD)
1178548342619-

Western Blot analysis of various cells using Histamine H1 Receptor Polyclonal Antibody diluted at 1:2000




+86-27-59760950

Immunofluorescence analysis of LOVO cells, using HRH1 Antibody. The picture on the right is blocked with the synthesized peptide.







Western blot analysis of lysates from COLO205 cells, using HRH1 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950