

## Histone deacetylase 10 rabbit pAb

Cat No.: ES2519

For research use only

## Overview

Product Name Histone deacetylase 10 rabbit pAb

Host species Rabbit

**Applications** WB;IHC;IF;ELISA

**Species Cross-Reactivity** Human; Mouse; Rat; Monkey **Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized peptide derived from human HDAC10. AA

range:10-59

**Specificity** Histone deacetylase 10 Polyclonal Antibody detects

endogenous levels of Histone deacetylase 10

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Histone deacetylase 10

Gene Name HDAC10

Cellular localizationCytoplasm . Nucleus . Excluded from nucleoli. .PurificationThe antibody was affinity-purified from rabbit<br/>antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 75kD
Human Gene ID 83933
Human Swiss-Prot Number Q969S8

Alternative Names HDAC10; Histone deacetylase 10; HD10

**Background** The protein encoded by this gene belongs to the

histone deacetylase family, members of which deacetylate lysine residues on the N-terminal part of

the core histones. Histone deacetylation modulates chromatin structure, and plays an important role in




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





transcriptional regulation, cell cycle progression, and developmental events. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011],

Western Blot analysis of various cells using Histone deacetylase 10 Polyclonal Antibody diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using HDAC10 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from COS7 cells, using HDAC10 Antibody. The lane on the right is blocked with the synthesized peptide.

