



# ICAD rabbit pAb

Cat No.:ES2589

For research use only


## Overview

<b>Product Name</b>	ICAD rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse;Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human DFFA. AA range:151-200
<b>Specificity</b>	ICAD Polyclonal Antibody detects endogenous levels of ICAD protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	DNA fragmentation factor subunit alpha
<b>Gene Name</b>	DFFA
<b>Cellular localization</b>	Cytoplasm.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	36kD
<b>Human Gene ID</b>	1676
<b>Human Swiss-Prot Number</b>	O00273
<b>Alternative Names</b>	DFFA; DFF1; DFF45; H13; DNA fragmentation factor subunit alpha; DNA fragmentation factor 45 kDa subunit; DFF-45; Inhibitor of CAD; ICAD
<b>Background</b>	Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied






by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using ICAD Polyclonal Antibody diluted at 1:1000





Western Blot analysis of 3T3 cells using ICAD Polyclonal Antibody diluted at 1:1000





Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using DFFA Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat, 293, and HeLa cells, using DFFA Antibody. The lane on the right is blocked with the synthesized peptide.

