

## IRF-3 rabbit pAb

## Cat No.:ES2643

For research use only

## Overview

Product Name	IRF-3 rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human;Mouse;Rat	
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.	
	Immunohistochemistry: 1/100 - 1/300. ELISA:	
	1/5000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
	peptide derived from human IRF3. AA	
	range:362-411	
Specificity	IRF-3 Polyclonal Antibody detects endogenous levels	
	of IRF-3 protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20 $^\circ\!{ m C}$ . Avoid repeated freeze-thaw cycles.	
Protein Name	Interferon regulatory factor 3	
Gene Name	IRF3	
<b>Cellular localization</b>	Cytoplasm . Nucleus . Mitochondrion . Shuttles	
	between cytoplasmic and nuclear compartments,	
	with export being the prevailing effect	
	(PubMed:10805757). When activated, IRF3	
	interaction with CREBBP prevents its export to the	
	cytoplasm (PubMed:10805757). Recruited to	
	mitochondria via TOMM70:HSP90AA1 upon Sendai	
	virus infection (PubMed:25609812)	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Observed band	48-55kd	
Human Gene ID	3661	
Human Swiss-Prot Number	Q14653	
		///////////////////////////////////////



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Alternative Names Background IRF3; Interferon regulatory factor 3; IRF-3 This gene encodes a member of the interferon regulatory transcription factor (IRF) family. The encoded protein is found in an inactive cytoplasmic form that upon serine/threonine phosphorylation forms a complex with CREBBP. This complex translocates to the nucleus and activates the transcription of interferons alpha and beta, as well as other interferon-induced genes. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2011],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C