

LRAT rabbit pAb

Cat No.:ES2724

For research use only

Overview

Product Name	LRAT rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
-	peptide derived from human LRAT. AA
	range:111-160
Specificity	LRAT Polyclonal Antibody detects endogenous levels
	of LRAT protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Lecithin retinol acyltransferase
Gene Name	LRAT
Cellular localization	Endoplasmic reticulum membrane ; Single-pass
	membrane protein . Rough endoplasmic reticulum .
	Endosome, multivesicular body . Cytoplasm,
	perinuclear region . Present in the rough
	endoplasmic reticulum and multivesicular body in
	hepatic stellate cells. Present in the rough
	endoplasmic reticulum and perinuclear region in
	endothelial cells (By similarity)
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	27kD
Human Gene ID	9227
5	



+86-27-59760950

ELKbio@ELKbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C

www.elkbiotech.com




Human Swiss-Prot Number Alternative Names

Background


095237

LRAT; Lecithin retinol acyltransferase; Phosphatidylcholine--retinol O-acyltransferase lecithin retinol acyltransferase (phosphatidylcholine--retinol O-acyltransferase)(LRAT) Homo sapiens The protein encoded by this gene localizes to the endoplasmic reticulum, where it catalyzes the esterification of all-trans-retinol into all-trans-retinyl ester. This reaction is an important step in vitamin A metabolism in the visual system. Mutations in this gene have been associated with early-onset severe retinal dystrophy and Leber congenital amaurosis 14. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2014],



Western Blot analysis of various cells using LRAT Polyclonal Antibody diluted at 1:500

Immunofluorescence analysis of HUVEC cells, using LRAT Antibody. The picture on the right is blocked with the synthesized peptide.





+86-27-59760950


ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Immunohistochemistry analysis of paraffin-embedded human prostate carcinoma tissue, using LRAT Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from COLO205 cells, using LRAT Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C