

## MARK2 rabbit pAb

## Cat No.:ES2747

For research use only

## Overview

Product Name	MARK2 rabbit pAb	
Host species	Rabbit	
Applications	WB;IF;ELISA	
Species Cross-Reactivity	Human;Mouse;Rat;Monkey	
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. Immunofluorescence	:
	1/200 - 1/1000. ELISA: 1/20000. Not yet tested in	
	other applications.	
Immunogen	The antiserum was produced against synthesized	
	peptide derived from human MARK2. AA	
	range:10-59	
Specificity	MARK2 Polyclonal Antibody detects endogenous	
	levels of MARK2 protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	Serine/threonine-protein kinase MARK2	
Gene Name	MARK2	
Cellular localization	Cell membrane; Peripheral membrane protein.	
	Cytoplasm. Lateral cell membrane. Cytoplasm,	
	cytoskeleton. Cell projection, dendrite . Cytoplasm .	
	Phosphorylation at Thr-596 by PRKCZ/aPKC and	
	subsequent interaction with 14-3-3 protein YWHAZ	
	promotes relocation from the cell membrane to the	
	cytoplasm.	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	85kD	
Human Gene ID	2011	
Human Swiss-Prot Number	Q7KZI7	
Alternative Names	MARK2; EMK1; Serine/threonine-protein kinase	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




Background

MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affinity-regulating kinase 2; PAR1 homolog; PAR1 homolog b; Par-1b; Par1b microtubule affinity regulating kinase 2(MARK2) Homo sapiens This gene encodes a member of the Par-1 family of serine/threonine protein kinases. The protein is an important regulator of cell polarity in epithelial and neuronal cells, and also controls the stability of microtubules through phosphorylation and inactivation of several microtubule-associating proteins. The protein localizes to cell membranes. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2009],

Western Blot analysis of various cells using MARK2 Polyclonal Antibody



Immunofluorescence analysis of A549 cells, using MARK2 Antibody. The picture on the right is blocked with the synthesized peptide.






+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C