

MNT rabbit pAb

Cat No.: ES2809

For research use only

Overview

Product Name MNT rabbit pAb

Host species Rabbit

WB;ELISA;IHC **Applications**

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 **Immunogen**

The antiserum was produced against synthesized

peptide derived from human MNT. AA range:315-364

Specificity MNT Polyclonal Antibody detects endogenous levels

of MNT protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Protein Name Max-binding protein MNT

Gene Name MNT **Cellular localization** Nucleus.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml Observed band 62kD **Human Gene ID** 4335 **Human Swiss-Prot Number** Q99583

Alternative Names MNT; BHLHD3; ROX; Max-binding protein MNT;

Class D basic helix-loop-helix protein 3; bHLHd3;

Myc antagonist MNT; Protein ROX

Background The Myc/Max/Mad network comprises a group of

transcription factors that co-interact to regulate


gene-specific transcriptional activation or

repression. This gene encodes a protein member of the Myc/Max/Mad network. This protein has a basic-Helix-Loop-Helix-zipper domain (bHLHzip) with







which it binds the canonical DNA sequence CANNTG, known as the E box, following heterodimerization with Max proteins. This protein is likely a transcriptional repressor and an antagonist of Myc-dependent transcriptional activation and cell growth. This protein represses transcription by binding to DNA binding proteins at its N-terminal Sin3-interaction domain. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using MNT Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western Blot analysis of A549 cells using MNT Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).







+86-27-59760950