

Mob1A rabbit pAb

Cat No.:ES2810

For research use only

Overview

Product Name Mob1A rabbit pAb

Host species Rabbit

WB;IHC;IF;ELISA **Applications**

Species Cross-Reactivity Human; Mouse; Monkey

Recommended dilutions Western Blot: 1/500 - 1/2000.

> Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

The antiserum was produced against synthesized **Immunogen**

peptide derived from human MOL1A. AA

range:101-150

Specificity Mob1A Polyclonal Antibody detects endogenous

levels of Mob1A protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

MOB kinase activator 1B **Protein Name**

Gene Name MOB1B

Cellular localization Cytoplasm . Nucleus .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 25kD **Human Gene ID** 92597 **Human Swiss-Prot Number** Q7L9L4

Alternative Names MOB1B; MOB4A; MOBKL1A; MOB kinase activator

1B; Mob1 homolog 1A; Mob1A; Mob1B; Mps one

binder kinase activator-like 1A

Background The protein encoded by this gene is similar to the

> yeast Mob1 protein. Yeast Mob1 binds Mps1p, a protein kinase essential for spindle pole body

duplication and mitotic checkpoint regulation. Three




+86-27-59760950 ELKbio@ELKbiotech.com



transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011],

Western Blot analysis of various cells using Mob1A Polyclonal Antibody



293 COS HUVEC 293
-- 117
-- 85
-- 48
-- 34
-- 26
-- 19
(kD)

Western blot analysis of lysates from 293, COS7, and HUVEC cells, treated with IFN 2500U/ml 30', using MOL1A Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

