

## NDUFB9 rabbit pAb

Cat No.: ES2910

For research use only

## Overview

Product Name NDUFB9 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human NDUFB9. AA

range:102-151

**Specificity** NDUFB9 Polyclonal Antibody detects endogenous

levels of NDUFB9 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Store at -20°C.** Avoid repeated freeze-thaw cycles.

Protein Name NADH dehydrogenase [ubiquinone] 1 beta

subcomplex subunit 9

Gene Name NDUFB9

**Cellular localization** Mitochondrion inner membrane; Peripheral

membrane protein; Matrix side.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 22kD
Human Gene ID 4715
Human Swiss-Prot Number Q9Y6M9

Alternative Names NDUFB9; LYRM3; UQOR22; NADH dehydrogenase

[ubiquinone] 1 beta subcomplex subunit 9; Complex

I-B22; CI-B22; LYR motif-containing protein 3; NADH-ubiquinone oxidoreductase B22 subunit


**Background** The protein encoded by this gene is a subunit of the






mitochondrial oxidative phosphorylation complex I (nicotinamide adenine dinucleotide: ubiquinone oxidoreductase). Complex I is localized to the inner mitochondrial membrane and functions to dehydrogenate nicotinamide adenine dinucleotide and to shuttle electrons to coenzyme Q. Complex I deficiency is the most common defect found in oxidative phosphorylation disorders and results in a range of conditions, including lethal neonatal disease, hypertrophic cardiomyopathy, liver disease, and adult-onset neurodegenerative disorders. Pseudogenes of this gene are found on chromosomes five, seven and eight. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2015],

Western Blot analysis of various cells using NDUFB9 Polyclonal Antibody diluted at 1:500




Western Blot analysis of 293 cells using NDUFB9 Polyclonal Antibody diluted at 1:500



+86-27-59760950








COLO COLO 293

Western blot analysis of lysates from COLO205 cells and 293 cells, using NDUFB9 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using NDUFB9 antibody.

