

Neuromedin-S rabbit pAb

Cat No.: ES2932

For research use only

Overview

Product Name Neuromedin-S rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human NMS. AA

range:104-153

Specificity Neuromedin-S Polyclonal Antibody detects

endogenous levels of Neuromedin-S protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Neuromedin-S

Gene Name NMS **Cellular localization** Secreted .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band24kDHuman Gene ID129521Human Swiss-Prot NumberQ5H8A3

Alternative Names NMS; Neuromedin-S

Background This gene encodes a member of the neuromedin

family of neuropeptides. The encoded preproprotein is proteolytically processed to generate a biologically

active neuropeptide that plays a role in the

regulation of circadian rhythm, anorexigenic action, antidiuretic action, cardiovascular function and




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




stimulation of oxytocin and vasopressin release. [provided by RefSeq, May 2016],

Western Blot analysis of various cells using Neuromedin-S Polyclonal Antibody diluted at 1:1000



MCF-7 293 HUVECHepG2 JK JK -- 117 -- 85 -- 48 -- 34 -- 26 -- 19 (kD)

Western blot analysis of lysates from Jurkat, HepG2, HUVEC, 293, and MCF-7 cells, using NMS Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

