

## HPA1 rabbit pAb

Cat No.: ES4083

For research use only

## Overview

Product Name HPA1 rabbit pAb

Host species Rabbit

**Applications** WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300.

ELISA: 1/20000. Not yet tested in other applications.

The antiserum was produced against synthesized.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

HPSE. AA range:241-290

**Specificity** HPA1 Polyclonal Antibody detects endogenous levels

of HPA1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Heparanase

Gene Name HPSE

**Cellular localization** Lysosome membrane; Peripheral membrane

protein. Secreted. Nucleus. Proheparanase is

secreted via vesicles of the Golgi. Interacts with cell membrane heparan sulfate proteoglycans (HSPGs).

Endocytosed and accumulates in endosomes.

Transferred to lysosomes where it is proteolytically cleaved to produce the active enzyme. Under certain stimuli, transferred to the cell surface. Associates with lipid rafts. Colocalizes with SDC1 in endosomal/lysosomal vesicles. Accumulates in perinuclear lysosomal vesicles. Heparin retains proheparanase in the extracellular medium (By

similarity). .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

**Clonality** Polyclonal



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




Concentration
Observed band
Human Gene ID
Human Swiss-Prot Number
Alternative Names

1 mg/ml 62kD 10855 Q9Y251


HPSE; HEP; HPA; HPA1; HPR1; HPSE1; HSE1; Heparanase; Endo-glucoronidase; Heparanase-1; Hpa1

Background

Heparan sulfate proteoglycans are major components of the basement membrane and extracellular matrix. The protein encoded by this gene is an enzyme that cleaves heparan sulfate proteoglycans to permit cell movement through remodeling of the extracellular matrix. In addition, this cleavage can release bioactive molecules from the extracellular matrix. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011],



Western Blot analysis of K562 cells using HPA1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100

