

## Macroglobulin $\alpha$ -2 rabbit pAb

## Cat No.:ES4098

For research use only

## Overview

Product Name	Macroglobulin α-2 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300.
	ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
C	peptide derived from the Internal region of human
	A2M. AA range:871-920
Specificity	Macroglobulin α-2 Polyclonal Antibody detects
	endogenous levels of Macroglobulin α-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Alpha-2-macroglobulin
Gene Name	A2M
Cellular localization	Secreted .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	170kD
Human Gene ID	2
Human Swiss-Prot Number	P01023
Alternative Names	A2M; CPAMD5; FWP007; Alpha-2-macroglobulin;
	Alpha-2-M; C3 and PZP-like alpha-2-macroglobulin
	domain-containing protein 5
Background	Alpha-2-macroglobulin is a protease inhibitor and
	cytokine transporter. It inhibits many proteases,
	including trypsin, thrombin and collagenase. A2M is
	implicated in Alzheimer disease (AD) due to its
	ability to mediate the clearance and degradation of




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C







A-beta, the major component of beta-amyloid deposits. [provided by RefSeq, Jul 2008],

Western Blot analysis of AD293 cells using Macroglobulin  $\alpha$ -2 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:100





+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C