

CD71/TfR rabbit pAb

Cat No.: ES4104

For research use only

Overview

Product Name CD71/TfR rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300.

ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the N-terminal region of

human TFRC. AA range:101-150

Specificity CD71 Polyclonal Antibody detects endogenous

levels of CD71 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Transferrin receptor protein 1

Gene Name TFRC

Cellular localization Cell membrane ; Single-pass type II membrane

protein. Melanosome. Identified by mass

spectrometry in melanosome fractions from stage I to stage IV. .; [Transferrin receptor protein 1, serum

form]: Secreted.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 85kD
Human Gene ID 7037
Human Swiss-Prot Number P02786

Alternative Names TFRC; Transferrin receptor protein 1; TR; TfR; TfR1;

Trfr; T9; p90; CD71

Background This gene encodes a cell surface receptor necessary

for cellular iron uptake by the process of




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeq, Sep 2015],


Western Blot analysis of HuvEc cells using CD71 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100



+86-27-59760950







Western blot analysis of lysate from HUVEC cells, using TFRC Antibody.

