

FMO3 rabbit pAb

Cat No.:ES4183

For research use only

Overview

Product Name	FMO3 rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human;Rat;Mouse;	
Recommended dilutions	Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300.	
	ELISA: 1/20000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
	peptide derived from the Internal region of human	
	FMO3. AA range:101-150	
Specificity	FMO3 Polyclonal Antibody detects endogenous	
. ,	levels of FMO3 protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	Dimethylaniline monooxygenase [N-oxide-forming]	
	3	
Gene Name	FMO3	
Cellular localization	Microsome membrane ; Single-pass membrane	
	protein . Endoplasmic reticulum membrane ;	
	Single-pass membrane protein .	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	58kD	
Human Gene ID	2328	
Human Swiss-Prot Number	P31513	
Alternative Names	FMO3; Dimethylaniline monooxygenase	
	[N-oxide-forming] 3; Dimethylaniline oxidase 3; FMO	
	II; FMO form 2; Hepatic flavin-containing	
	monooxygenase 3; FMO 3; Trimethylamine	
	monooxygenase	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

flavin containing monooxygenase 3(FMO3) Homo sapiens Flavin-containing monooxygenases (FMO) are an important class of drug-metabolizing enzymes that catalyze the NADPH-dependent oxygenation of various nitrogen-, sulfur-, and phosphorous-containing xenobiotics such as therapeutic drugs, dietary compounds, pesticides, and other foreign compounds. The human FMO gene family is composed of 5 genes and multiple pseudogenes. FMO members have distinct developmental- and tissue-specific expression patterns. The expression of this FMO3 gene, the major FMO expressed in adult liver, can vary up to 20-fold between individuals. This inter-individual variation in FMO3 expression levels is likely to have significant effects on the rate at which xenobiotics are metabolised and, therefore, is of considerable interest to the pharmaceutical industry. This transmembrane protein localizes to the endoplasmic reticulum of many tissues. Alternative splicing of this gen



Western Blot analysis of HeLa cells using FMO3 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C