

IL-18R rabbit pAb

Cat No.: ES4202

For research use only

Overview

Product Name IL-18R rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

IL18R1. AA range:231-280

Specificity IL-18R Polyclonal Antibody detects endogenous

levels of IL-18R protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Interleukin-18 receptor 1

Gene Name IL18R1

Cellular localization Membrane ; Single-pass type I membrane protein . **Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 63kD
Human Gene ID 8809
Human Swiss-Prot Number Q13478

Alternative Names IL18R1; IL1RRP; Interleukin-18 receptor 1; IL-18R-1;

IL-18R1; CD218 antigen-like family member A; CDw218a; IL1 receptor-related protein; IL-1Rrp;

IL1R-rp; CD218a

Background The protein encoded by this gene is a cytokine

receptor that belongs to the interleukin 1 receptor family. This receptor specifically binds interleukin 18

(IL18), and is essential for IL18 mediated signal




+86-27-59760950 ELKbio@ELKbiotech.com




transduction. IFN-alpha and IL12 are reported to induce the expression of this receptor in NK and T cells. This gene along with four other members of the interleukin 1 receptor family, including IL1R2, IL1R1, ILRL2 (IL-1Rrp2), and IL1RL1 (T1/ST2), form a gene cluster on chromosome 2q. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2013],

Western Blot analysis of 293 cells using IL-18R Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from 293 cells, using IL18R1 Antibody.



+86-27-59760950

