

## FGF-4 rabbit pAb

## Cat No.:ES4282

For research use only

## Overview

Product Name	FGF-4 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
-	peptide derived from the C-terminal region of
	human FGF4. AA range:151-200
Specificity	FGF-4 Polyclonal Antibody detects endogenous
	levels of FGF-4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Fibroblast growth factor 4
Gene Name	FGF4
Cellular localization	Secreted .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	22kD
Human Gene ID	2249
Human Swiss-Prot Number	P08620
Alternative Names	FGF4; HST; HSTF1; KS3; Fibroblast growth factor 4;
	FGF-4; Heparin secretory-transforming protein 1;
	HST; HST-1; HSTF-1; Heparin-binding growth factor
	4; HBGF-4; Transforming protein KS3
Background	The protein encoded by this gene is a member of
	the fibroblast growth factor (FGF) family. FGF family
	members possess broad mitogenic and cell survival
	activities and are involved in a variety of biological



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This gene was identified by its oncogenic transforming activity. This gene and FGF3, another oncogenic growth factor, are located closely on chromosome 11. Co-amplification of both genes was found in various kinds of human tumors. Studies on the mouse homolog suggested a function in bone morphogenesis and limb development through the sonic hedgehog (SHH) signaling pathway. [provided by RefSeq, Jul 2008],

Western Blot analysis of NIH-3T3, Hela cells using FGF-4 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C