

## Gβ 5 rabbit pAb

Cat No.: ES4534

For research use only

## Overview

**Product Name** Gβ 5 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human GNB5. AA

range:151-200

Specificity Gβ 5 Polyclonal Antibody detects endogenous levels

of Gβ 5 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Guanine nucleotide-binding protein subunit beta-5

Gene Name GNB5

Cellular localization Membrane.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band40kDHuman Gene ID10681Human Swiss-Prot Number014775

Alternative Names GNB5; Guanine nucleotide-binding protein subunit

beta-5; Gbeta5; Transducin beta chain 5

**Background** Heterotrimeric guanine nucleotide-binding proteins

(G proteins), which integrate signals between


receptors and effector proteins, are composed of an alpha, a beta, and a gamma subunit. These subunits are encoded by families of related genes. This gene



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





encodes a beta subunit. Beta subunits are important regulators of alpha subunits, as well as of certain signal transduction receptors and effectors.

Alternatively spliced transcript variants encoding different isoforms exist. [provided by RefSeq, Jul 2008],

Western Blot analysis of HepG2 cells using G $\beta$  5 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



+86-27-59760950