

FRS2 rabbit pAb

Cat No.:ES4559

For research use only

Overview

Product Name	FRS2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
<u> </u>	peptide derived from human FRS2. AA
	range:402-451
Specificity	FRS2 Polyclonal Antibody detects endogenous levels
	of FRS2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Fibroblast growth factor receptor substrate 2
Gene Name	FRS2
Cellular localization	Endomembrane system. Cytoplasmic,
	membrane-bound.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	85kD
Human Gene ID	10818
Human Swiss-Prot Number	Q8WU20
Alternative Names	FRS2; Fibroblast growth factor receptor substrate 2;
	FGFR substrate 2; FGFR-signaling adaptor SNT;
	Suc1-associated neurotrophic factor target 1; SNT-1
Background	function:Adapter protein that links FGR and NGF
C	receptors to downstream signaling pathways.
	Involved in the activation of MAP kinases.
	Modulates signaling via SHC1 by competing for a



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



common binding site on

NTRK1.,PTM:Phosphorylated on tyrosine residues upon stimulation by NGF.,PTM:Ubiquitinated when tyrosine phosphorylated and in a complex with GRB2. The unphosphorylated form is not subject to ubiquitination.,sequence caution:Translated as stop.,similarity:Contains 1 IRS-type PTB domain.,subcellular location:Cytoplasmic, membrane-bound.,subunit:Part of a complex containing FRS2, GRB2 and SOS1. Part of a complex containing GRB2 and CBL. Binds RET (By similarity). Binds FGFR1, SUC1, NTRK1, NTRK2, NTRK3 and SRC. The tyrosine-phosphorylated protein binds the SH2 domains of GRB2 and PTPN11.,tissue specificity:Highly expressed in heart, brain, spleen, lung, liver, skeletal muscle, kidney and testis.,

Western Blot analysis of HUVEC cells using FRS2 Polyclonal Antibody





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C