

Olfactory receptor 52E2 rabbit pAb

Cat No.: ES4675

For research use only

Overview

Product Name Olfactory receptor 52E2 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR52E2. AA

range:276-325

Specificity Olfactory receptor 52E2 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 52E2

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 52E2

Gene Name OR52E2

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 37kD
Human Gene ID 119678
Human Swiss-Prot Number O8NGJ4

Alternative Names OR52E2; Olfactory receptor 52E2

Background Olfactory receptors interact with odorant molecules


in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from




+86-27-59760950 ELKbio@ELKbiotech.com



single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],




Western Blot analysis of Jurkat cells using Olfactory receptor 52E2 Polyclonal Antibody



Western blot analysis of lysates from Jurkat cells, using OR52E2 Antibody. The lane on the right is blocked with the synthesized peptide.







Western blot analysis of the lysates from HeLa cells using OR52E2 antibody.

