



Olfactory receptor 10T2 rabbit pAb

Cat No.:ES4732

For research use only


Overview

Product Name	Olfactory receptor 10T2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human OR10T2. AA range:231-280
Specificity	Olfactory receptor 10T2 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10T2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 10T2
Gene Name	OR10T2
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	30kD
Human Gene ID	128360
Human Swiss-Prot Number	Q8NGX3
Alternative Names	OR10T2; Olfactory receptor 10T2; Olfactory receptor OR1-3
Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from






single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western Blot analysis of 293T-UV cells using Olfactory receptor 10T2 Polyclonal Antibody diluted at 1:500



Western blot analysis of lysates from HUVEC and COLO cells, using OR10T2 Antibody. The lane on the right is blocked with the synthesized peptide.

