

Casein Kinase Iα (phospho Thr321) rabbit

pAb

Cat No.:ES4856

For research use only

Overview

Product Name	Casein Kinase Iα (phospho Thr321) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human Casein Kinase I alpha
	around the phosphorylation site of Tyr321. AA
	range:287-336
Specificity	Phospho-Casein Kinase Iα (Y321) Polyclonal
	Antibody detects endogenous levels of Casein
	Kinase I $lpha$ protein only when phosphorylated at
	Y321.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Casein kinase I isoform alpha
Gene Name	CSNK1A1
Cellular localization	Cytoplasm . Cytoplasm, cytoskeleton, microtubule
	organizing center, centrosome . Chromosome,
	centromere, kinetochore . Nucleus speckle .
	Cytoplasm, cytoskeleton, cilium basal body .
	Cytoplasm, cytoskeleton, spindle . Localizes to the
	centrosome in interphase cells, and to kinetochore
	fibers during mitosis. Also recruited to the keratin
	cytoskeleton (PubMed:23902688)
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
+86-27-59760950	ELKbio@ELKbiotech.com www.elkbiotech.com





23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Concentration	1 mg/ml
Observed band	35kD
Human Gene ID	1452
Human Swiss-Prot Number	P48729
Alternative Names	CSNK1A1; Casein kinase I isoform alpha; CKI-alpha; CK1
Background	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. It can phosphorylate a large number of proteins. Participates in Wnt signaling. Phosphorylates CTNNB1 on 'Ser-45'.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CK1 Ser/Thr protein kinase family. Casein kinase I subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Monomer. Interacts with the Axin
Alternative Names	CSNK1A1; Casein kinase I isoform alpha; CKI-alp CK1 catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Casein kinases are operationally defined by their preferential utiliz of acidic proteins such as caseins as substrates. can phosphorylate a large number of proteins. Participates in Wnt signaling. Phosphorylates CTNNB1 on 'Ser-45'.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to protein kinase superfamily. CK1 Ser/Thr protein kinase family. Casein kinase I subfamily.,similarity:Contains 1 protein kinase

Western Blot analysis of VEC cells using Phospho-Casein Kinase Ia (Y321) Polyclonal Antibody





Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Casein Kinase I alpha (Phospho-Tyr321) Antibody




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Western blot analysis of lysates from HUVEC cells, using Casein Kinase I alpha (Phospho-Tyr321) Antibody. The lane on the right is blocked with the phospho peptide.

Immunohistochemical analysis of paraffin-embedded human spleen tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C