

Cystatin SA rabbit pAb

Cat No.:ES4874

For research use only

Overview

Product Name	Cystatin SA rabbit pAb
Host species	Rabbit
Applications	IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human CST2. AA range:11-60
Specificity	Cystatin SA Polyclonal Antibody detects endogenous
	levels of Cystatin SA protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cystatin-SA
Gene Name	CST2
Cellular localization	Secreted .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	
Human Gene ID	1470
Human Swiss-Prot Number	P09228
Alternative Names	CST2; Cystatin-SA; Cystatin-2; Cystatin-S5
Background	The cystatin superfamily encompasses proteins that
	contain multiple cystatin-like sequences. Some of
	the members are active cysteine protease inhibitors,
	while others have lost or perhaps never acquired
	this inhibitory activity. There are three inhibitory
	families in the superfamily, including the type 1
	cystatins (stefins), type 2 cystatins and the
	kininogens. The type 2 cystatin proteins are a class



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a secreted thiol protease inhibitor found at high levels in saliva, tears and seminal plasma. [provided by RefSeq, Jul 2008],

Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using CST2 Antibody. The picture on the right is blocked with the synthesized peptide.





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C